



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2011

---

## **Notes on the occurrence of *Syngnathus rostellatus* (Teleostei, Syngnathidae) in the Mediterranean**

Hablützel, Pascal I ; Wilson, Anthony B

**Abstract:** *Syngnathus rostellatus* is a nearshore pipefish species whose distributional range extends along the European Atlantic coast between Bergen (NO) and the Bay of Biscay (ES). Several recent articles suggest that this species has experienced a major range expansion of more than 4000 km into the eastern Mediterranean, but a critical review of these studies indicates that the majority of these reports are based on specimen misidentifications. Considering a reliable report of *S. rostellatus* from the Mediterranean coast near Gibraltar, it appears that the current distribution of this species is restricted to the north-eastern Atlantic Ocean and the southern Mediterranean coast of the Iberian Peninsula.

DOI: <https://doi.org/10.1017/S1755267211000558>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-60899>

Journal Article

Accepted Version

Originally published at:

Hablützel, Pascal I; Wilson, Anthony B (2011). Notes on the occurrence of *Syngnathus rostellatus* (Teleostei, Syngnathidae) in the Mediterranean. *Marine Biodiversity Records*, 4:1-4.

DOI: <https://doi.org/10.1017/S1755267211000558>

1 Accepted Manuscript - *Marine Biodiversity Records*

2

3 **Notes on the occurrence of *Syngnathus rostellatus* (Teleostei, Syngnathidae) in the**  
4 **Mediterranean**

5

6 Pascal I. Hablützel<sup>a1 a2 c1</sup> and Anthony B. Wilson<sup>a1</sup>

7

8 <sup>a1</sup> Institute of Evolutionary Biology and Environmental Studies, University of Zurich,  
9 Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

10

11 <sup>a2</sup> Laboratory of Animal Diversity and Systematics, K.U. Leuven, Charles Deberiotstraat  
12 32, B-3000 Leuven, Belgium

13

14 <sup>c1</sup> Correspondence should be addressed to: P.I. Hablützel, Laboratory of Animal Diversity  
15 and Systematics, K.U. Leuven, Charles Deberiotstraat 32, B-3000 Leuven, Belgium email:  
16 pascal.habluetzel@bio.kuleuven.ch

17 Abstract

18 *Syngnathus rostellatus* Nilsson, 1855 is a nearshore pipefish species whose distributional  
19 range extends along the European Atlantic coast between Bergen (NO) and the Bay of  
20 Biscay (ES). Several recent articles suggest that this species has experienced a major  
21 range expansion of more than 4000 km into the eastern Mediterranean, but a critical  
22 review of these studies indicates that the majority of these reports are based on specimen  
23 misidentifications. Considering a reliable report of *S. rostellatus* from the Mediterranean  
24 coast near Gibraltar, it appears that the current distribution of this species is restricted to  
25 the northeastern Atlantic Ocean and the southern Mediterranean coast of the Iberian  
26 peninsula.

27

28 Keywords: range expansion, species identification, *Syngnathus rostellatus*, pipefish

## 29 INTRODUCTION

30 There is increasing evidence that human-mediated environmental changes, including the  
31 erosion of geographical barriers, the introduction of exotic species, and habitat  
32 degradation, are influencing the distribution of species (e.g. Rahel, 2007; Hiddink & ter  
33 Hofstede, 2008). The accelerated rate of human modifications of the natural environment is  
34 paralleled by an increasing number of reports documenting extensions and contractions of  
35 the historical ranges of many species (e.g. Azzurro, 2008; Sax & Gaines, 2003; Thomas,  
36 2010). The value of these studies is entirely dependent on the accurate identification of  
37 specimens, and the misidentification of material can lead to erroneous conclusions of  
38 major shifts in species' ranges.

39

40 Close to 40 nominal species of the globally-distributed genus *Syngnathus* are currently  
41 recognized in the Eschmeyer's Catalog of Fishes (2010). Of these, eleven are known to  
42 occur in Europe, making this the most species-rich group of syngnathid fishes in the region  
43 (Dawson, 1986). While nine of these species names are in common use (e.g. Dawson,  
44 1986), two species from the Black Sea (*S. affinis*, Eichwald, 1831 and *S. argentatus*, Pallas,  
45 1814) were only recently reinstated by Kuiter (2009). Kuiter (2009) provides no  
46 justification for the reinstatement of these species names, and a more systematic analysis of  
47 Black Sea pipefish will be necessary in order to clarify their species status. The taxonomic  
48 status of at least two additional species is also unclear. The nearshore *S. agassiz*  
49 Michahelles, 1829 sensu Canestrini (1872) was synonymized with *S. abaster* by Lueken  
50 (1967), but was this species was reinstated by Roig (1979) on the basis of an exhaustive  
51 analysis of historical and contemporary collections from the Balearic Islands and Spanish

52 coast – this Spanish-language report appears to have been overlooked by subsequent  
53 researchers. Similarly, genetic and morphometric analyses of pipefish material from the  
54 Mediterranean and Black Seas suggest that *S. nigrolineatus* Eichwald, 1831 may also  
55 represent a distinct species (Hablützel & Wilson, unpublished data). Clearly, the taxonomy  
56 of *Syngnathus* pipefish is in a state of flux, and the genus is in need of a full systematic  
57 revision.

58

59 Of the nine commonly-recognized species, two (*S. phlegon* Risso, 1827 and *S. schmidtii*  
60 Popov, 1928) are pelagic, while the others (*S. abaster* Risso, 1827 sensu Canestrini (1872),  
61 *S. acus* Linnaeus, 1758, *S. rostellatus* Nilsson, 1855, *S. taenionotus* Canestrini, 1871, *S.*  
62 *tenuirostris* Rathke, 1837, *S. typhle* Linnaeus, 1758 and *S. variegatus* Pallas, 1811) are all  
63 nearshore inhabitants (Dawson, 1986), and are dominant members of nearshore eelgrass  
64 habitats along the European coastline.

65

66 *S. rostellatus* (Nilsson's pipefish) was originally described from western Sweden (Nilsson,  
67 1855) and the northern French *S. dumerilii* Moreau, 1870 in Duméril (1870) is now  
68 considered a junior synonym of this species (Fries *et al.*, 1895; Wheeler, 1973).

69 Ehrenbaum (1905-1909) defined the distribution range of *S. rostellatus* as the Atlantic  
70 coast between Bergen to the Sea of Biscay (Figure 1). *S. rostellatus* is found at moderate  
71 densities in the Kattegat of western Sweden, and while there are scattered reports of  
72 individual specimens in the southern Baltic (e.g. Ehrenbaum, 1905-1909; Otterstøm,  
73 1917), a recent exhaustive survey of this region failed to detect this species (HELCOM  
74 2007). These earlier descriptions have been complemented by subsequent reports of *S.*

*rostellatus* from the northern Irish coast (Douglas and Egan 1983) and sites as far south as Málaga, Spain (Helling (1943); Reina-Hervás *et al.* (1981-1982)) (Figure 1). *S. rostellatus* has recently been reported from sites within the Mediterranean as far east as southern Turkey (Louisy, 2002; Gökoglu *et al.*, 2004; Ben Amor *et al.*, 2008) (Figure 1), suggesting that this species may be experiencing a major range extension.

Here, we provide a morphological diagnosis of *S. rostellatus*, and use this tool to evaluate recent report of *S. rostellatus* outside its historical distributional range (Dawson, 1986). Our analyses indicate that the majority of these recent reports stem from the misidentification of specimens and that the distribution of this species remains restricted to the northeastern Atlantic Ocean and the Mediterranean coast of the Iberian peninsula.

## MATERIALS AND METHODS

Pipefish individuals were identified on the basis of the meristic and morphometric data provided in the original articles and, if available, from pictures. The original description of *S. rostellatus* by Nilsson (1855), together with subsequent descriptions by Moreau in Duméril (1870) and Dawson (1986) were used as references, and these descriptions were supplemented with the analysis of ethanol-preserved specimens from Northern Spain (Sada) and Western Sweden (Fiskebäckskil) (Table 1). Methods for the measurements and meristic counts used in this article follow Dawson (1986).

## RESULTS

### Identification

*Syngnathus rostellatus* differs from all other European species of the genus by the following combination of characters: distal margins of body rings without spine-like points; 13-17 trunk rings; 35-42 tail rings; 32-45 dorsal fin rays; 10-13 pectoral fin rays; lack of brown dots in the dorsal fin; and small size ( $< 17$  cm TL) (Table 1). While the meristic counts of the Black Sea population of *S. abaster* (*S. abaster nigrolineatus* sensu Berg (1949)) overlap with those of *S. rostellatus*, this species can be clearly distinguished from *S. rostellatus* both morphologically and genetically (Hablützel & Wilson, unpublished data)).

## **Distribution**

Almeida (1986) reported 12 *S. rostellatus* specimens (including eight pregnant males) from Vila Nova de Milfontes (PT) (Figure 1). The reproductive status of the individual specimens was not indicated in the original paper, but the data provided indicate that the smallest pregnant male in this collection had a TL of  $\leq 12.1$  cm. *Syngnathus rostellatus*, *S. abaster*, *S. agassiz* and *S. typhle* are the only four coast-associated species of *Syngnathus* in Western Europe which are reported to reach sexual maturity at this size. *Syngnathus abaster* has fewer dorsal fin rays ( $\leq 31$ ), *S. agassiz* has less tail rings ( $\leq 35$ ) and *S. typhle* has more pectoral fin rays ( $\geq 14$ ) than does *S. rostellatus* (Table 1). The meristic counts of the specimens collected by Almeida (1986) are thus consistent with the identification of these specimens as *S. rostellatus* (Table 1). Almeida (1986) also re-examined specimens from Praia de Mira (PT) (Figure 1) collected by Helling (1943) and identified these individuals as *S. rostellatus*. On the basis of the details provided in Almeida (1986), we conclude that all of these individuals are indeed *S. rostellatus*.

121

122 Reports of *S. rostellatus* from Málaga (Figure 1) by Lozano y Rey (1919) and Reina-  
123 Hervás *et al.* (1981-1982) also appear to be reliable. While details on the specimens  
124 analyzed by Lozano y Rey (1919) were not provided in the original publication, Hervás *et*  
125 *al.* (1981-1982) provided detailed information on five specimens collected from the same  
126 region. The low number of trunk rings detected in these specimens (16-17; Table 1) clearly  
127 separates them from juvenile *S. acus* (trunk rings: 18-19) which are otherwise similar in  
128 their general appearance. The *S. rostellatus* from Málaga also differ from *S. abaster*, *S.*  
129 *agassiz* and *S. typhle* in meristic counts (see above).

130

131 In contrast to these reliable reports of *S. rostellatus* from outside its historical range,  
132 several recent descriptions of *S. rostellatus* from sites in Tunisia and Turkey (Figure 1)  
133 appear to be in error. Photographs of the individuals included in both of these studies are  
134 provided in the original articles (Gökoglu *et al.*, 2004; Ben Amor *et al.*, 2008). Both  
135 pictures show large (18.7 cm and 21.1 cm TL), long-snouted species with high numbers of  
136 trunk rings (18 for the Tunisian and > 17 for the Anatolian individual (note that Gökoglu *et*  
137 *al.* used a non-standard trunk counting method which differs from other publications of  
138 syngnathid pipefish (e.g. Dawson, 1986))), indicating that neither of these specimens are *S.*  
139 *rostellatus* (Table 1; Figure 2). The specimens illustrated by Gökoglu *et al.* (2004) and Ben  
140 Amor *et al.* (2008) most closely resemble either *S. tenuirostris* or *S. acus* (Figure 2),  
141 species which are known to occur in the Mediterranean region (Kaup, 1856; Dawson,  
142 1986).

143



Louisy (2002) reported a single *S. rostellatus* female from Banyuls-sur-Mer, France (Figure 1). The author did not provide meristic data, but included two color pictures of the living specimen. The low number of dorsal fin rays (29 or 30; Table 1) and the flattened form of the snout (versus the slightly convex snout of *S. rostellatus* (Figure 2)) suggest that the specimen is actually *S. abaster*. The report of *S. taenionotus* from Southern France in Louisy (2002) also appears to be in error, and is based on an apparent misidentification of a *S. typhle* individual (data not shown). Unfortunately, the photographs and identifications used by Louisy (2002) have been incorporated into a recent global survey of syngnathid species (Kuitert 2009), further propagating these errors.

Confusion about the distributional range of *S. rostellatus* also exists at its northeastern periphery in the Southern Baltic Sea. Kuitert (2009) recently reported two specimens from Wismar, Germany. We argue that both pictured specimens are misidentified and in fact are *S. typhle*, recognizable by the elevation of the snout evident in these individuals, diagnostic for this species (see Figure 2). One of the two specimens also shows a color pattern of white spots which is known only from juvenile *S. typhle* and is not found in *S. rostellatus*.

#### **The lasting impact of historical errors on the European biogeography of *Syngnathus***

Dawson's (1986) key to European syngnathid species has been an important reference for researchers working on European *Syngnathus* species. Unfortunately, an error in the illustration included in this text has perpetuated confusion relating to species-level identification. While the meristic counts provided in Dawson (1986) are accurate, the illustration of *S. rostellatus* included in this report was derived from a sketch originally

published in Fries *et al.* (1895) and subsequently reprinted by Poll (1947) in his review of Belgian marine fishes. Unfortunately, while the original illustration (Fries *et al.* 1895) referred to *S. typhle* (plate XXIX, figure 1), Poll (1947) mislabeled this specimen as *S. rostellatus* in both the text and the figure legend of his book (p. 186, Figure 126), and Dawson kept this description in his text.

Poll (1947) made a second labeling error in his *Fauna de Belgique*, including an illustration of *S. rostellatus* originally published in Fries *et al.* (1895) in his description of *S. typhle* (plate XXVIII, figure 8b; note that this specimen is also inconsistently labeled as both *S. rostellatus* and *S. acus* in the original publication). Such labeling errors have undoubtedly contributed to the confusion surrounding the identification of European *Syngnathus* species, and we have attempted to remedy this situation here, including photographs of all of commonly recognized nearshore *Syngnathus* species known to occur in this region (Figure 2).

## CONCLUSIONS

Considering the reliable reports of *S. rostellatus* from the Portuguese and the Spanish Mediterranean coasts, we suggest that the current range of *S. rostellatus* is broader than that indicated by Dawson (1986). The contemporary range of this species extends from the Norwegian coast as far south as the western Mediterranean at Málaga, Spain. This distributional pattern coincides with the present frontal system between Almeria and Oran, with separates Atlantic waters to the west from Mediterranean waters to the east (Tintore *et al.*, 1988). Reports of *S. rostellatus* from Mediterranean sites east of Málaga appear to be

erroneous, and stem from specimen misidentifications.

## ACKNOWLEDGMENTS

We thank Alexandra Wegmann and Jasmin Winkler for their assistance in species identification. Filip Volckaert and two anonymous reviewers provided helpful comments on an earlier draft of this article. We further thank the Senckenberg Museum Frankfurt for access to their pipefish collection. The study was supported by the University of Zurich, Switzerland.

## REFERENCES

- Almeida A.J.** (1986) Presence des poissons *Syngnathus rostellatus* (Syngnathidae), *Clinitrachus argentatus* (Clinidae) et *Sphoeroides spencleri* (Tetraodontidae) sur la côte du Portugal. *Ciência Biológica: Ecology and Systematics (Portugal)* 6 (1-2), 1 - 7.
- Azzurro E.** (2008) The advance of thermophilic fishes in the Mediterranean Sea: overview and methodological questions. In CIESM *Climate warming and related changes in Mediterranean marine biota*. N° 35 in *CIESM Workshop Monographs* [F. Briand, Ed.]. Monaco: CIESM, pp. 39-45.
- Ben Amor M.M., Ben Salem M., Ben Soussi J. and Capapé C.** (2008) Occurrence of Nilsson's Pipefish *Syngnathus rostellatus* (Osteichthyes: Syngnathidae) in the Gulf of Tunis (Central Mediterranean). *Vie et Milieu – Life and Environment* 58 (3-4), 189 - 192.

213 **Berg L.S.** (1949) Fresh-water fishes of Soviet Union and adjacent countries. III. Freshw.  
 214 Fish. USSR, Guide Fauna USSR. no. 30, 927 - 1382. (in Russian, English translation  
 215 available, 1965, Smithsonian Institution, Washington D.C., by Israel Program of Scientific  
 216 Translation)  
 217

218 **Canestrini G.** (1872) *Fauna d'Italia - Parte terza - Pesci*. Milano: Dottor Francesco  
 219 Vallardi.  
 220

221 **Dawson C.E.** (1986) Syngnathidae. In Whitehead P.J.P., Bauchot M.-L., Hureau J.-C.,  
 222 Nielson J. and Tortonese E. (eds) *Fishes of the Northeastern Atlantic and the*  
 223 *Mediterranean, Vol II*. Paris: UNESCO, pp. 628 - 639.  
 224

225 **Douglas D.J. and Egan D.** (1983) The distribution of *Syngnathus rostellatus* Nilsson  
 226 (Pisces: Syngnathidae) in Irish coastal waters. *Irish Naturalists Journal* 21 (4), 173 - 175.  
 227

228 **Duméril A.H.A.** (1870) *Histoire Naturelle des Poissons ou Ichthyologie Générale. Tome*  
 229 *Second. Ganoïdes, Dipnés, Lophobranches*. Paris: Librairie Encyclopédique de Roret.  
 230

231 **Ehrenbaum E.** (1905-1909) I. Eier und Larven von Fischen. In Ehrenbaum E. (ed)  
 232 *Nordisches Plankton - Zoologischer Teil - Erster Band - Eier und Larven von Fischen,*  
 233 *andere Eier und Cystem*. Kiel und Leipzig: Verlag von Lipsius & Tischer, pp. 1 - 414.  
 234

235 **Eschmeyer W.N.** (ed.) (2010) *Catalog of Fishes*. Electronic version (25 October 2010).

236 <http://research.calacademy.org/ichthyology/catalog/fishcatmain.asp>  
 237

238 **Fries B., Ekström C.U. and C.J. Sundevall** (1895) *A History of Scandinavian Fishes.*  
 239 *Part II.* 2nd ed., revised and completed by Professor F. A. Smitt. Stockholm: P. A. Norstedt  
 240 & Söner.  
 241

242 **Gökoglu M., Bodur T. and Kaya Y.** (2004) First records of *Hippocampus fuscus* and  
 243 *Syngnathus rostellatus* (Osteichthyes: Syngnathidae) from the Anatolian coast  
 244 (Mediterranean Sea). *Journal of the Marine Biological Association of the United Kingdom*  
 245 84 (5), 1093 - 1094.  
 246

247 **HELCOM** (2007) HELCOM Red list of threatened and declining species of lampreys and  
 248 fish of the Baltic Sea. *Baltic Sea Environmental Proceedings* 109, 1 - 40 pp.  
 249

250 **Helling H.** (1943) Novo catálogo de peixes de Portugal em colecção no Museu de  
 251 Zoologia da Universidade de Coimbra. *Memórias e Estudos do Museu Zoológico da*  
 252 *Universidade de Coimbra* s. 1 (149), 1 - 110.  
 253

254 **Hiddink J.G. and ter Hofstede R.** (2008) Climate induced increases in species richness of  
 255 marine fishes. *Global Change Biology* 14, 453 - 460.  
 256

257 **Kaup J.J.** (1856) *Catalogue of lophobranchiate fish in the collection of the British*  
 258 *Museum.* London: British Museum.

259

260 **Kuiter, R.H.** (2009) *Seahorses and their relatives*. Seaford: Aquatic Photographics.

261

262 **Louisy P.** (2002) *Meeresfische: Westeuropa, Mittelmeer*. Stuttgart: Verlag Eugen Ulmer.

263

264 **Lozano y Rey L.** (1919) Los Peces de la Fauna Ibérica en la Colección del Museo en 1 de  
265 enero de 1919. *Trabajos del Museo Nacional de Ciencias Naturales Seria Zoológica* 39, 1  
266 - 112.

267

268 **Lueken W.** (1967) Süßwasserfische der Türkei - 5. Teil Syngnathidae. *Mitteilungen*  
269 *Hamburgisches Zoologisches Museum und Institut* 64, 127 - 146.

270

271 **Nilsson S.** (1855) *Skandinavisk fauna. Fjerde Delen: Fiskarna*. Lund: Berlingska  
272 Boktryckeriet.

273

274 **Otterstøm C.V.** (1917) *Danmarks Fauna - Fisk III. Fastkæbede, Buskgællede, Ganoider,*  
275 *Tværmunde og Rundmunde*. Copenhagen: G.E.C. Gads Forlag.

276

277 **Poll M.** (1947) *Faune de Belgique: Poissons marins*. Brussels: Musée royal d'Histoire  
278 naturelle de Belgique.

279

280 **Rahel F.J.** (2007) Biogeographic barriers, connectivity and homogenization of freshwater  
281 faunas: it's a small world after all. *Freshwater Biology* 52 (4), 696 - 710.

282

283 **Reina-Hervás J.A., Muñoz R. and Blasco M.** (1981-1982) Presencia de teleosteos  
284 atlánticos en el Mediterráneo occidental. *Monografías y Trabajos de Zoología*  
285 (Universidad de Málaga) 3-4, 49 - 56.

286

287 **Roig A.** (1979) *Singnathus agassizi* Kaup, Canestrini: nombre válido para unos singnátidos  
288 capturados en Sa Rápita (Mallorca-Islas Baleares) (Pisces, Syngnathidae). *Miscelánea*  
289 *Zoológica* (Barcelona) 5, 83 - 92.

290

291 **Sax D.F. and Gaines S.D.** (2003) Species diversity: from global decreases to local  
292 increases. *Trends in Ecology and Evolution* 18 (11), 561 - 566.

293

294 **Thomas C.D.** (2010) Climate, climate change and range boundaries. *Diversity and*  
295 *Distribution* 16, 488 - 495.

296

297 **Tintore J, Violette P.E., Blade I. and Cruzado A.** (1988) A study of an intense density  
298 front in the eastern alboran sea: the Almeria-Oran front. *Journal of Physical Oceanography*  
299 18, 1384 - 1397.

300

301 **Wheeler A.** (1973) Syngnathidae. In Hureau J.C., and Monod T. (eds) *Check-list of the*  
302 *fishes of the north-eastern Atlantic and of the Mediterranean*. Paris: UNESCO, pp. 274 -  
303 279.

304 **Fig. 1.** The traditional distribution of *S. rostellatus* according to Dawson (1986) (black),  
305 extended by reports from the Irish (Douglas & Egan 1983), Portuguese (Almeida, 1986)  
306 and Spanish (Reina-Hervás *et al.*, 1981-1982) coasts and the southern Baltic Sea  
307 (Ehrenbaum, 1905-1909; Otterstøm, 1917) (grey); collection localities of *S. rostellatus*  
308 discussed in the text are indicated (circles), as are reports of *S. rostellatus* based on  
309 apparent misidentifications of specimens (stars).

310

311 **Fig. 2.** Photographs of the head morphology of the nearshore *Syngnathus* species discussed  
312 in this article, along with an indication of the collection locality of the individual  
313 specimens. **A:** *S. abaster*, Scardovari (IT); **B:** *S. acus*, Tasende (ES); **C:** *S. agassiz*, Naples  
314 (IT); **D:** *S. rostellatus*, Sada (ES); **E:** *S. taenionotus*, Scardovari (IT); **F:** *S. tenuirostris*,  
315 Crimea (UA); **G:** *S. typhle*, Venice (IT); **H:** *S. variegatus*, Crimea (UA). Scale bar equals 1  
316 cm. Pictures (A-E, G) taken from specimens collected by the authors and collaborators  
317 (A,B,D,E,G) or from the collection of the Senckenberg Museum Frankfurt (C; SMF8334).  
318 Drawings (F, H) after Rathke (1837), see Kuitert (2009) for photographs of these species.




Figure 1



Figure 2

319



**Table 1.** Meristic characters of *S. rostellatus* from the original species description (Nilsson, 1855) and from more recent studies of the species. Morphological data for specimens outside the expected range for *S. rostellatus* are indicated in bold.

Location	N	Trunk rings	Tail rings	Dorsal fin rays	Pectoral fin rays	Length of largest specimen [cm TL]	Conclusion	Reference
Sweden	n.a.	16	40	32-34	10	12.4-14.8 (5-6 tum)	<i>S. rostellatus</i>	Nilsson, 1855
Côte du Hâvre (FR)	n.a.	14	36-39	34-36	11-12	10.3	<i>S. rostellatus</i>	Moreau in Duméril, 1870
Europe	n.a.	13-17	37-42	33-45	10-13	17.0	<i>S. rostellatus</i>	Dawson, 1986
Sada (ES)	30	14-16	39-42	37-42	10-12	13.0	<i>S. rostellatus</i>	this study
Fiskebäckskil (SE)	9	14-15	39-41	38-43	11-12	13.7	<i>S. rostellatus</i>	this study
Vila Nova de Milfontes (PT)	12	14-16	35-42	34-40	10-11	14.0	<i>S. rostellatus</i>	Almeida, 1986
Praia de Mira (PT)	6	15	39-42	37-42	11	16.9	<i>S. rostellatus</i>	Almeida, 1986
Málaga (ES)	5	16-17	34-41	36-38	11-12	n.a.	<i>S. rostellatus</i>	Reina-Hervás <i>et al.</i> , 1981-1982
Gulf of Tunis (TN)	1	<b>18</b>	39	35	12	<b>21.1</b>	<i>Syngnathus</i> sp.	Ben Amor <i>et al.</i> , 2008
Anatolian Coast (TR)	1	<b>&gt; 17</b>	41	33	13	<b>18.7</b>	<i>Syngnathus</i> sp.	Gökoglu <i>et al.</i> , 2004
Banyuls-sur-Mer (FR)	1	n.a.	n.a.	<b>29 or 30</b>	n.a.	n.a.	<i>S. abaster</i>	Louisy, 2002
Wismar (DE)	2	n.a.	n.a.	n.a.	n.a.	n.a.	<i>S. typhle</i>	Kuiter, 2009

N, number; 1 tum = 2.47 cm